

Preoperative Instructions for Ablation

Procedure:	Date:	Check- in Time	:
Intended Discharge Plan: You m change at the time of your proce	,	, ,	•
Hospital:	Address:		
Check-In Location: Hospital Mai	n Floor Entrance – Admissions	5	
<u>Plan Ahead</u> :			
Same day discharges will not li	kely leave the hospital until la	te afternoon or early eve	ning.
• For 24 hours, you must have so	omeone stay with you and you	ı cannot drive.	
• Plan for someone to drive you	home from the hospital. Ride	services (Taxi, Uber, Lyft)	are not allowed.
The Day Before Your Procedure	<u>::</u>		
• Please have nothing to eat or o	drink after midnight the night l	before the procedure, inc	luding gum and candy.
 You may take your prescription our office. 	n medications the morning of	the procedure unless inst	ructed otherwise by
Medication Specifics:			
• Last pre-procedure dose of you	ur blood thinner	will be on date:	in the am/pm.
• Do not take your Jardiance, Fa	rxiga for 3 days before proced	lure.	
Do not take your water pill	the morn	ing of your procedure.	
• Do not take any vitamins or mi	nerals the morning of your pro	ocedure.	
Diabetic patients ONLY			
• Do not take your usual oral dia	betic medications the morning	g of the procedure.	
• Take 1/2 of your insulin the mo	orning of the procedure.		

Bring to the Hospital:

- Bring an accurate list of your medications including the medication name, pill strength, dose you take, and times of the day you take it
- Bring your CPAP or BiPAP, if applicable
- Bring any inhalers you will need

What symptoms are expected after my ablation?

- You may experience mild chest discomfort that worsens with deep breaths or lying flat. This should gradually improve.
- You may have occasional palpitations that last seconds or minutes related to inflammation.
- It is normal to have some groin site soreness, a small lump under the skin, and bruising that can even extend down the leg.

What are my activity restrictions after ablation?

- Groin Site Restrictions: no lifting more than 10 lbs, straining during bowel movements, excessive bending
 over, stooping, walking long distances, running, or climbing many flights of stairs for 1 week.
- *SVT/AVN/PVC/VT/Typical Flutter:* Groin site restrictions (above) during 1st week, then no restrictions thereafter.
- A-fib/Cardioneural/SVT with transeptal puncture: In addition to groin site restrictions during the first week, be a "couch potato" for the first few days. 1 week after ablation, you may begin to ease back into your daily routine (without exercise). Continue to avoid heavy lifting for 2 weeks.. weeks post-procedure, you may slowly ramp-up your exercise intensity over the following 2 weeks. No strenuous exercise or scuba diving for 1 month. No activity restrictions 1-month post-procedure.

When can I return to work?

- *SVT/AVN/PVC/VT/Typical Flutter:* Return to sedentary job in a few days. Wait 1 week to return if work involves heavy physical exertion.
- A-fib/Cardioneural/SVT with transeptal puncture: Return to sedentary job in 1 week. Wait 2 weeks to return if work involves heavy physical exertion.

When can I travel?

- PVC/A-fib: local patients should avoid travel for 2 weeks post-procedurally.
- All other ablation types: local patients should avoid travel for 1 week post-procedurally.

When should I follow-up?

- A-fib/Cardioneural/SVT with transeptal puncture or any ablation by Dr. Alyesh: 1 week visit with EP RN
- All Ablations: 1 month visit in Arrhythmia Clinic. Subsequent MD visit timing is procedure-specific.

When can I have other procedures, dental cleanings, or surgeries?

- A-fib/PVC: none for 3 months (sometimes PVC ablations only need to wait 1 month)
- All other ablation types: none for 1 month